

Magnetar Driven Bubbles and the Origin of Collimated Outflows in Gamma-ray Bursts

N. Bucciantini^{1*}, E. Quataert¹, J. Arons^{1,2,3}, B. D. Metzger^{1,2}, T. A. Thompson⁴

¹*Astronomy Department and Theoretical Astrophysics Center,*

University of California at Berkeley, 601 Campbell Hall, Berkeley CA, 94720, USA

²*Physics Department, University of California at Berkeley*

³*Kavli Institute for Particle Astrophysics and Cosmology, Stanford University*

⁴*Lyman Spitzer Jr. Fellow, Department of Astrophysical Sciences, Peyton Hall, Ivy Lane, Princeton, NJ, 08544, USA*

Accepted . Received ; in original form

ABSTRACT

We model the interaction between the wind from a newly formed rapidly rotating magnetar and the surrounding supernova shock and host star. The dynamics is modeled using the two-dimensional, axisymmetric thin-shell equations. In the first $\sim 10 - 100$ seconds after core collapse the magnetar inflates a bubble of plasma and magnetic fields behind the supernova shock. The bubble expands asymmetrically because of the pinching effect of the toroidal magnetic field, even if the host star is spherically symmetric, just as in the analogous problem of the evolution of pulsar wind nebulae. The degree of asymmetry depends on $E_{\text{mag}}/E_{\text{tot}}$, the ratio of the magnetic energy to the total energy in the bubble. The correct value of $E_{\text{mag}}/E_{\text{tot}}$ is uncertain because of uncertainties in the conversion of magnetic energy into kinetic energy at large radii in relativistic winds; we argue, however, that bubbles inflated by newly formed magnetars are likely to be significantly more magnetized than their pulsar counterparts. We show that for a ratio of magnetic to total power supplied by the central magnetar $\dot{E}_{\text{mag}}/\dot{E}_{\text{tot}} \lesssim 0.1$ the bubble expands relatively spherically. For $\dot{E}_{\text{mag}}/\dot{E}_{\text{tot}} \gtrsim 0.3$, however, most of the pressure in the bubble is exerted close to the rotation axis, driving a collimated outflow out through the host star. This can account for the collimation inferred from observations of long-duration gamma-ray bursts (GRBs). Outflows from magnetars become increasingly magnetically dominated at late times, due to the decrease in neutrino-driven mass loss as the young neutron star cools. We thus suggest that the magnetar-driven bubble initially expands relatively spherically, enhancing the energy of the associated supernova, while at late times it becomes progressively more collimated, producing the GRB. The same physical processes may operate in more modestly rotating neutron stars to produce asymmetric supernovae and lower energy transients such as X-ray flashes.

Key words: Stars: neutron; stars: supernovae: general; gamma-rays: bursts; stars: winds, outflows; magnetic field; MHD

1 INTRODUCTION

In the first few seconds after core collapse in a massive star, a proto neutron star (PNS) cools and contracts on its Kelvin-Helmholtz cooling timescale ($\sim 10 - 100$ s), radiating its gravitational binding energy ($\sim 10^{53}$ ergs) in neutrinos (Burrows 1986, Pons et al. 1999). The cooling epoch is accompanied by mass loss driven by neutrino energy deposi-

tion in the atmosphere of the PNS (e.g. Duncan et al. 1986, Qian & Woosley 1996, Thompson et al. 2001).

A subset of neutron stars are inferred to have magnetic field strengths of $\sim 10^{14} - 10^{15}$ G (“magnetars”; see Woods & Thompson 2004 for a review). If some magnetars are born with millisecond rotation periods (e.g., Duncan & Thompson 1992, Thompson & Duncan 1993), the combination of rapid rotation and strong magnetic fields makes the winds from young magnetars significantly more energetic than the thermally driven winds from slowly rotating neutron stars. In addition, as the neutrino-driven mass

* E-mail: nbucciantini@astro.berkeley.edu

loss decreases during the Kelvin-Helmholtz cooling epoch, the wind from a proto-magnetar becomes increasingly magnetically-dominated and the flow eventually becomes relativistic. For this reason, proto-magnetars have been considered as a possible central engine for long-duration gamma ray bursts (GRBs)¹ and hyper-energetic supernovae (Usov 1992, Thompson 1994, Wheeler et al. 2000, Thompson et al. 2004), and as a possible source of ultra-high energy cosmic rays (Blasi et al. 2000, Arons 2003).

The discovery that GRBs are at cosmological distances confirmed that the isotropic energy scale for the gamma-ray emission from GRBs is $\sim 10^{52} - 10^{53}$ ergs (see, e.g., Woosley & Bloom 2006). However, the interpretation of afterglow observations (“jet breaks”) suggested that GRBs are powered by collimated jets and that the intrinsic energy in relativistic material is $\sim 10^{51}$ ergs (e.g., Frail et al. 2001). This interpretation has become less clear in recent years because of the complex time-dependence in SWIFT X-ray afterglow observations and the lack of evidence for X-ray jet breaks in the first ~ 10 days (e.g., Sato et al. 2007, Burrows & Racusin 2007). Nonetheless, the case for collimated outflows from GRBs is compelling. Theoretically, the association of many long-duration GRBs with supernovae (Woosley & Bloom 2006) sets the natural energy scale for GRBs at $\sim 10^{51} - 10^{52}$ ergs. In addition, estimates of the energy in relativistic outflows in GRBs from late time radio observations provide lower limits of the same order, although the true energy could in principle be much higher (see, e.g., Eichler & Waxman 2005).

In the collapsar model (e.g., MacFadyen & Woosley 1999), the collimated outflows from GRBs are accounted for by jets produced by an accretion flow onto a central black hole. In the magnetar model, the origin of such collimated outflows is less clear. Relativistic magnetized outflows by themselves do not efficiently self-collimate (e.g., Lyubarsky & Eichler 2001). Although observations of pulsar wind nebulae (PWNe) – which are intrinsically far more relativistic than GRBs – show jet-like features (e.g., Weisskopf et al. 2000, Pavlov et al. 2001, Gaensler et al. 2002), these are believed to be only mildly relativistic outflows produced by the interaction between the pulsar wind and the surrounding expanding supernova (SN) remnant (Komissarov & Lyubarsky 2004, Del Zanna et al. 2004). In this paper, we explore the hypothesis that collimated outflows from newly formed magnetars can likewise be produced by the interaction between the magnetar wind and the surrounding host star.

Our physical picture is that the fast trans-magnetosonic magnetar wind shocks on the relatively slow outgoing SN envelope, creating a subsonic bubble of plasma and magnetic fields inside its host star. Because of the strong toroidal magnetic field and the accompanying pinch, an anisotropic pressure distribution between the pole and equator is set up within the cavity defined by the SN shock and the incoming magnetar wind. For simplicity we assume that (1) an outgoing SN shock has created a central evacuated cavity and (2) the surrounding host star is spherically symmetric. Assumption (1) allows us to model the problem of

interest as a free magnetar wind interacting with the expanding envelope created by a SN shock that is in turn sweeping through the host star. Spectral modeling of the hyper-energetic supernovae associated with several GRBs suggests massive progenitor stars (e.g., Iwamoto et al. 1998; Mazzali et al. 2006). This has been interpreted as indicating that GRBs are associated with the formation of black holes. However, there is increasing evidence that some Galactic magnetars arise from massive stars with ZAMS masses of $\approx 40M_{\odot}$ (e.g., Muno et al. 2006). Thus our assumption of a successful core-collapse SN leaving behind a rapidly rotating magnetar is quite reasonable given current observational constraints on the progenitors of magnetars and GRBs. Our assumption (2) that the host star is spherically symmetric may be conservative. Multi-dimensional simulations of core-collapse in the presence of rapid rotation and strong poloidal magnetic fields find that the explosion may occur preferentially along the rotation axis (e.g., LeBlanc & Wilson 1970, Burrows et al. 2007). It is presumably easier to produce a late-time collimated outflow in this case, since a low pressure, low inertia channel has already been created.

A full magnetohydrodynamic (MHD) simulation of the interaction between a magnetar wind and its host star would require resolving a very wide radial dynamic range. In addition, the physical conditions in the wind at large distances – in particular, the magnetization of the wind – are not fully understood (§2). For these reasons, we believe that it is fruitful to solve a model problem that allows one to readily explore the parameter space of magnetar-host star interactions – the thin-shell approximation provides such a model.

In the thin-shell approximation, one assumes that the material swept-up by the wind from the central object is confined to a geometrically thin shell, whose dynamics is then evolved (e.g., Giuliani 1982). This model has been extensively applied in the study of the interaction of stellar winds with their surrounding environment, both in the case of momentum driven winds (see, e.g., Canto 1980; Canto et al. 1996; Wilkin 2000) and in the case of pressure driven winds (e.g., Chevalier & Luo 1994). The evolution of magnetized PWNe bounded by an expanding SN remnant (Begelman & Li 1992) is the closest analogue to the problem we consider in this paper. In a number of cases, more detailed numerical simulations have confirmed the validity of the thin-shell model (see, e.g., Stevens et al. 1992; Bucciantini 2002 for hydrodynamical examples). Most importantly for our purposes, axisymmetric relativistic MHD simulations by van der Swaluw (2003) and Del Zanna et al. (2004) have shown that the overall shape of PWNe resembles that predicted by the thin-shell model of Begelman & Li (1992). For these reasons we believe that the thin-shell shell approximation is a useful tool for studying the structure and evolution of bubbles formed by magnetar winds inside their progenitor stars. In addition, these calculations can define the most interesting parameter space for future relativistic MHD simulations.

The remainder of this paper is organized as follows. In § 2 we discuss the general properties of proto-magnetar winds, and how they evolve in the ~ 100 seconds after core-collapse. We also discuss the equilibrium structure of the magnetized bubble created by the magnetar wind behind the SN shock. Section 3 summarizes the thin-shell equations. In § 4 we present our results for the evolution of the SN shock

¹ In what follows we typically drop the phrase “long-duration” for conciseness and refer to long-duration GRBs simply as GRBs.




Figure 1. Evolution of a magnetar wind as a function of time since core bounce, based on the evolutionary calculations of Metzger et al. (2007). *Top:* Energy loss rate *Bottom:* Magnetization at the light cylinder. Models are for an initial period of $P = 1$ ms, $B = 10^{15}$ G (dot dashed), $B = 3 \times 10^{15}$ G (solid), and $B = 10^{16}$ G (dotted) and an initial period of $P = 2$ ms and $B = 10^{16}$ G (dashed).

due to the asymmetric pressure produced by the interior magnetized bubble. In § 5 we summarize our conclusions and discuss the implications of our results for understanding observations of long-duration gamma-ray bursts, X-ray flashes, and asymmetric supernovae. In the Appendix we present self-similar solutions that provide insight into how the shape of the bubble is related to its magnetization and the conditions in the ambient medium.

2 PROTOMAGNETAR EVOLUTION AND BUBBLE STRUCTURE

2.1 Protomagnetar Wind Evolution

In order to set the stage for the thin-shell evolutionary calculations that follow, this section summarizes some of the key properties of outflows from young rapidly rotating magnetars. Metzger et al. (2007) (hereafter M07) solved the one-dimensional (monopolar) neutrino-heated non-relativistic

MHD wind problem for conditions appropriate to young magnetars. These calculations provide the mass-loss rate ($\dot{M}$) and energy-loss rate ($\dot{E}$) from the magnetar as a function of parameters such as the neutrino luminosity, magnetic field strength B , and rotation rate Ω . The calculation of $\dot{M}$ is applicable even if the wind is relativistic because the mass-loss is set in the non-relativistic subsonic portion of the wind at small radii. The calculations of M07 include the neutrino micro-physics necessary for direct application to PNSs. Their calculations were, however, restricted to monopolar field structure. A complementary set of calculations was carried out by Bucciantini et al. (2006) (hereafter B06), who studied aligned dipolar (axisymmetric) non-relativistic and relativistic MHD winds from neutron stars assuming an adiabatic equation of state. M07 showed that their results could be combined with those of B06 to provide evolutionary models for early magnetar spin-down, including the transition from non-relativistic to relativistic outflows as the neutrino-driven mass loss decreases.

Figure 1 shows the energy loss rate $\dot{E}$ and magnetization σ_{LC} as a function of time from the evolutionary calculations of M07 for several combinations of (dipole) magnetic field strengths and magnetar birth period. The values of $B \approx 10^{15} - 10^{16}$ G and $P \approx 1 - 2$ ms are chosen to be characteristic of PNSs that produce conditions suitable for producing GRBs or hyper-energetic supernovae. The magnetization in Figure 1 is defined by

$$\sigma_{LC} \equiv \Phi_B^2 \Omega^2 / \dot{M} c^3, \quad (1)$$

where Φ_B is the total open magnetic flux per 4π steradian (Michel 1969), $\dot{M}$ is the mass loss rate, and σ_{LC} is evaluated at the light cylinder. Winds with $\sigma_{LC} \lesssim 1$ are non-relativistic while those with $\sigma_{LC} \gtrsim 1$ are relativistic. The calculations in Figure 1 assume that the PNS luminosity decreases in time $\propto t^{-1}$ until $t = 40$ s, motivated by the cooling evolution of Pons et al. (1999). We note, however, that the cooling of strongly magnetized rapidly rotating PNSs is not well understood, which is a source of uncertainty in the determination of $\dot{M}(t)$, $\dot{E}(t)$ and $\sigma_{LC}(t)$.

The energy loss rates $\dot{E}$ in Figure 1 are a factor of ~ 10 larger than the “vacuum dipole” rate for the given value of Ω and B . There are two reasons for this. First, when the wind is non-relativistic ($\sigma_{LC} \lesssim 1$), the energy loss rate is larger than in the relativistic limit, with $\dot{E} \propto \dot{M}^{1/3} (\dot{M}^{3/5})$ for non-relativistic magnetized winds with a monopole (dipole) magnetic field structure (e.g., Thompson et al. 2004). In addition, the large mass loss rates accompanying PNS spin-down lead to excess open magnetic flux which enhances the spin-down of the PNS (see the simulations of B06). This is true even when the wind is moderately relativistic ($\sigma_{LC} \gtrsim 1$). The large energy loss rates shown in Figure 1 are sufficient to lead to appreciable spin-down of the PNS during the Kelvin-Helmholtz epoch. For example, for the model with $P = 1$ ms and $B = 3 \times 10^{15}$ G in Figure 1 (solid line), the PNS loses $\approx 80\%$ of its rotational energy in the first 40 seconds. This efficient spin-down is largely responsible for the fact that $\dot{E}$ decreases in time as the PNS cools (see Figure 1).²

² Two additional effects contribute to the decline in $\dot{E}$ with time. First, as the PNS cools, the mass loss rate $\dot{M}$ decreases. In the non-relativistic limit, the energy loss rate is proportional

As the PNS cools, the neutrino-driven mass loss decreases in time. This in turn causes a transition from a non-relativistic to relativistic wind, as shown explicitly in the plot of $\sigma_{LC}(t)$ in Figure 1. These calculations of σ_{LC} are based on equatorial spin-down models (M07), which probably underestimate the angle-averaged σ in the wind by a factor of few (B06). Nonetheless, the evolution from a moderately mass-loaded marginally relativistic wind ($\sigma_{LC} \sim 1$) to a highly magnetized Poynting flux dominated outflow ($\sigma_{LC} \gg 1$) is expected to be generic for cooling magnetars.

As we show in the next section, the impact of the magnetar on its host star depends critically on the strength of the magnetic field in the bubble created by the magnetar wind; the generation rate of the field in the bubble is in turn determined by the magnetization σ of the wind at large radii. In non-relativistic winds, the magnetic energy and kinetic energy are in approximate equipartition at large radii, with $E_{mag} \approx 2E_{kin}$ (e.g., Lamers & Cassinelli 1999). One-dimensional models of ideal relativistic winds, however, find that the asymptotic Lorentz factor of the wind is $\gamma_\infty \approx \sigma_{LC}^{1/3}$ and the asymptotic magnetization is $\sigma \approx \sigma_{LC}^{2/3}$ (Michel 1969, Goldreich & Julian 1970) so that most of the energy remains in the magnetic field at large radii. These results apply in the limit of $\sigma_{LC} \gg 1$. Relativistic MHD simulations (Bucciantini et al. 2007) show that for intermediate values of $\sigma_{LC} \lesssim 20$, a reasonable fraction of the magnetic energy is converted into kinetic energy at large distances, with rough equipartition obtaining by $\sim 10^4$ stellar radii.

In the limit of very high σ_{LC} , studies of PWNe (e.g. the Crab Pulsar) find that the wind must have low $\sigma \sim 10^{-2}$ at large radii (e.g., Kennel & Coroniti 1984, Begelman & Li 1992). Although there is no consensus on the mechanism responsible for the inferred decrease in pulsar wind magnetization at large radii, a prominent class of models relies on magnetic dissipation in the relativistic outflow over a large radial distance (e.g., Coroniti 1990; Lyubarsky & Kirk 2001; Kirk & Skjæraasen 2003). The physical conditions in proto-magnetar winds are quite different from those in pulsar winds (e.g., they are much denser so that there is no charge starvation). In addition, the distance to the termination shock is much smaller in the SN confined winds from young magnetars, ~ 10 light cylinder radii (see below) compared to more than 10^4 light cylinder radii in PWNe and in pulsar-Be star binaries. The reduced flow time between the light cylinder and the termination shock may mean that dissipation of magnetic energy in young magnetar winds is less complete than in pulsar winds. As a result, we suspect that the rate of injection of magnetic energy into bubbles created by protomagnetars may be significantly larger than that inferred in the PWNe context. Given the uncertainties, however, we treat the magnetization in the outflow, expressed as the ratio of the magnetic energy injection to the total power ($\dot{E}_{mag}/\dot{E}_{tot}$), as a free parameter in this paper, bearing in mind the generic evolution from $\sigma_{LC} \sim 1$ to $\sigma_{LC} \gg 1$ in Figure 1.

to $\dot{M}^{0.3-0.6}$ and thus decreases as well (this is relevant for the $P = 1$ ms, $B = 3 \times 10^{15}$ (solid line) and $P = 1$ ms, $B = 10^{15}$ G (dot-dashed) models in Figure 1 at early times). The decreasing mass loss rate also decreases the fraction of open magnetic flux and thus $\dot{E}$.

The models shown in Figure 1 assume that the wind from the central magnetar is freely expanding into a cavity evacuated by the outgoing SN shock. Formally, this requires that the radius of the fast magnetosonic point must be smaller than the radius of the SN shock; the latter is $R_s \sim 10^9$ cm in the first few seconds, which is indeed larger than the typical distance to the fast surface of $\sim 10 - 40$ neutron star radii (B06, for a millisecond rotator). As the freely expanding wind moves out, it interacts with the surrounding SN shock and previously shocked wind material. More precisely, the wind will reach a termination shock at which its kinetic energy is thermalized and the magnetic field is compressed. A correct determination of the size of the termination shock requires a full MHD model of the wind-bubble interaction (e.g., Del Zanna et al. 2004). As a rough guide to the relevant scales, however, we note that in the simple case of a constant $\dot{M}$ and $\dot{E}$, $\sigma_{LC} \sim 1$ wind moving into a spherically symmetric bubble, the termination shock is located at a radius $R_t \sim R_s(R_s/ct)^{1/2} \sim 0.1R_s \sim 10^8$ cm where t is the time since the onset of the wind (in sec). For $R_t < R < R_s$, the wind develops into a bubble of plasma and magnetic field confined by the SN shock and host star.

2.2 The Bubble Structure

If one neglects plasma flow inside the bubble, a simple solution for the structure inside the bubble ($R_t < R < R_s$) can be obtained in the case of the predominantly toroidal magnetic field expected at large radii in the wind. This solution was found by Begelman & Li (1992). We reproduce several of its features here because they are important to our model.

The Begelman & Li (1992) solution will be valid as long as typical flow speeds do not exceed the local sound speed. In the case of a relativistic magnetized bubble the sound speed ranges from $c/\sqrt{3}$ to c . It is possible that, close to the termination shock, post shock flow can move with high velocities (Del Zanna et al. 2004), but in the bulk of the bubble, typical speeds are expected to be a small fraction of c , unless the cavity itself expands at a considerable fraction of the speed of light. Indeed, as long as the expansion velocity of the shell is small compared to the sound speed inside the bubble, the plasma inside will always relax to pressure equilibrium, independent of the energy distribution in the wind (be it primarily polar as for a non-relativistic wind or primarily equatorial as for a relativistic wind). Neglecting the fluid flow, the structure is given by the assumption of magnetohydrostatic equilibrium. Assuming axisymmetry, the momentum equations become:

$$\frac{\partial}{\partial z} \left(p + \frac{B^2}{8\pi} \right) = 0, \quad \frac{\partial}{\partial r} \left(p + \frac{B^2}{8\pi} \right) + \frac{B^2}{4\pi r} = 0, \quad (2)$$

where r is the cylindrical radius, p is the pressure, and B the toroidal magnetic field in the bubble. The first equation simply states that isobaric surfaces are coaxial cylinders.

If entropy is constant along each flow streamline in the bubble then the continuity equation can be written as:

$$\frac{\partial}{\partial r} (p^{1/\Gamma} r v_r) + \frac{\partial}{\partial z} (p^{1/\Gamma} r v_z) = 0. \quad (3)$$

where Γ is the adiabatic index of the fluid. Comparing this with the flux-freezing condition for the toroidal magnetic field yields

$$p \propto (B/r)^\Gamma. \quad (4)$$

For the case of a relativistic plasma ($\Gamma = 4/3$), equation (4) can be used in the r -momentum equation to find

$$p = \frac{p_n}{\zeta^2}, \quad \frac{B^2}{8\pi} = \frac{9p_n r^2}{16\zeta^3 H^2}, \quad (5)$$

where ζ is the solution of the following equation:

$$\left(\zeta + \frac{9r^2}{32H^2}\right)^2 - \zeta^3 = 0. \quad (6)$$

The solution for the pressure in the bubble given by equations (2)-(6) depends on two parameters. One of these, the pressure on the axis p_n , determines the overall magnitude of the pressure in the bubble. The other, the scale height H of the pressure distribution, determines the pressure stratification in the bubble. In Figure 2 we plot the normalized pressure profile derived from the solution of the above equations. The total pressure is higher along the axis ($r = 0$) and asymptotically decreases as r^{-2} . The region close to the axis contains a low σ plasma and is essentially pressure dominated, while at larger distances the plasma is magnetically dominated, and the ratio of magnetic to thermal pressure increases linearly with the distance. Equipartition is reached for $r/H \sim 2$. The results in Figure 2 assume a relativistic plasma with $\Gamma = 4/3$, which corresponds to $\sigma_{LC} \gtrsim 1$ in Figure 1. The magnetar wind may be non-relativistic at very early times, so that $\Gamma = 5/3$ is more appropriate. For $\Gamma = 5/3$ the pressure profiles are qualitatively similar to those in Figure 2, although the scale height H is a factor of ≈ 2 smaller for a given ratio of magnetic to total energy in the bubble. For simplicity, we simply set $\Gamma = 4/3$ in all of our calculations.

The scale height H and the asymmetry of the pressure distribution can be expressed in terms of the ratio of the magnetic energy to total energy in the bubble. To quantify this effect, consider a spherical bubble of radius R and total energy E . The pressure along the axis is given by

$$p_n \approx 8 \times 10^{22} \bar{P} \left(\frac{E}{10^{51} \text{ ergs}} \right) \left(\frac{R}{10^9 \text{ cm}} \right)^{-3} \text{ ergs cm}^{-3}. \quad (7)$$

The dimensionless number $\bar{P}$ is the pressure on the axis relative to that in an unmagnetized bubble. Figure 3 shows $\bar{P}$ and the scale height H/R as a function of $E_{\text{mag}}/E_{\text{tot}}$, the ratio of the magnetic to total energy in the bubble (similar results are obtained for the self-similar solutions described in the Appendix; see, e.g., Figure A2). Magnetized bubbles have $\bar{P} \gg 1$ and $H \ll R$ (where R is the radius of the bubble, not the cylindrical radius within the bubble used above and in Figure 2). Figure 2 shows that, due to the pinching effect of the toroidal magnetic field, the pressure in the bubble in this case will be concentrated along the axis and so the bubble will expand asymmetrically. By contrast, very weakly magnetized bubbles have $H \gtrsim R$ and roughly constant pressure throughout. Note that a magnetization of $E_{\text{mag}}/E_{\text{tot}} \gtrsim 0.1$ is required to make $H \lesssim R$ and the pressure distribution in the bubble relatively asymmetric.

We now calculate how the swept-up shell in the host star responds to the pressure produced by the magnetized bubble created by the central magnetar.




Figure 2. Pressure structure for a relativistic plasma containing toroidal magnetic field in magneto-hydrostatic equilibrium (based on Begelman & Li 1992). The solid line (A) is the total pressure, normalized to the value on the axis. The dotted line (B) is the magnetic pressure, normalized to the total pressure on the axis. The dashed line (C) is the ratio of the magnetic to the total pressure, while the dash-dotted line (D) is the ratio of the magnetic to the thermal pressure. Bubbles with weak magnetic fields have large values of H relative to the size of the bubble (see Figure 3) and thus only the $r \ll H$ part of this plot is applicable: as a result the pressure is relative uniform and the system will expand spherically. By contrast, bubbles with appreciable magnetic fields have smaller values of H and thus the pressure on the axis is significantly larger than the pressure near the equator. Such bubbles will expand asymmetrically.




Figure 3. Dimensionless pressure $\bar{P}$ (see eq. [7]) and scale-height H as a function of the magnetic energy in the bubble, for the case of spherical bubble. $\bar{P}$ is the ratio of the pressure on axis to the pressure in an unmagnetized spherical bubble. For even modest magnetic energy, the pressure distribution becomes highly anisotropic with $\bar{P} \gg 1$ and $H \ll R$.

3 THE THIN-SHELL EQUATIONS

The equations describing the evolution of an axisymmetric bubble can be greatly simplified if one assumes that the swept-up material is confined in a thin-shell, so that the dynamics can be described in the “thin-shell approximation.” This approximation is valid as long as the thickness of the shell is small compared to the radius of curvature. The thin-shell equations account for conservation of mass and momentum. A detailed derivation of the equations can be found in Giuliani (1982) where corrections for the thickness of the shell are also discussed. In the case of infinitesimally thin shell they are given by:

$$\tan \xi = -\frac{1}{R} \frac{\partial R}{\partial \theta}, \quad (8)$$

$$u_{\perp} = \cos \xi \frac{\partial R}{\partial t}, \quad (9)$$

$$u_{\parallel} = \sin \xi \frac{\partial R}{\partial t}, \quad (10)$$

$$\frac{\partial(A\sigma)}{\partial t} = -\rho_o(v_{\perp o} - u_{\perp})A + \rho_i(v_{\perp i} - u_{\perp})A - \frac{\partial}{\partial \theta} [R \sin \theta \sigma (v_{\parallel} - u_{\parallel})], \quad (11)$$

$$\frac{\partial(A\sigma \mathbf{v})}{\partial t} = -[\rho_o(v_{\perp o} - u_{\perp})\mathbf{v}_o + \mathbf{e}_{\perp}(p_o + B_o^2/8\pi)]A + [\rho_i(v_{\perp i} - u_{\perp})\mathbf{v}_i + \mathbf{e}_{\perp}(p_i + B_i^2/8\pi)]A - \frac{\partial}{\partial \theta} [R \sin \theta \sigma (v_{\parallel} - u_{\parallel})\mathbf{v}], \quad (12)$$

$$A = \left(\frac{R^2 \sin \theta}{\cos \xi} \right). \quad (13)$$

where ξ is the angle between the radial direction and the normal to the shell surface, A is the effective area of each element of the shell, and σ is the surface density. The suffixes $\perp$ and $\parallel$ represent the directions perpendicular and parallel to the shell (and they are time dependent), while i and o label the conditions at the inner and outer edge of the shell. The velocity $v_{\perp} = u_{\perp}$ is the expansion velocity of the shell perpendicular to itself, $u_{\parallel}$ is the expansion velocity parallel to itself, and $v_{\parallel}$ is the flow velocity along the shell.

Equation (11) represents conservation of mass along the shell, while equation (12) describes momentum conservation in the shell. Both equations include a flux of the relevant quantity along the shell itself, and source terms due to the inner and outer media. As discussed in Giuliani (1982), these equations employ a Lagrangian remapping along the shell, and can be applied only as long as $R(\theta)$ is a single valued function of θ .

The evolution of the thin shell depends on the force driving it (“i”) and on the conditions in the external medium (“o”). The solution in §2.2 describes the inner conditions used in this study. In many cases of interest, the outer medium is sufficiently cold and the magnetic field is sufficiently weak that their contribution to the pressure term in equation (12) can be neglected. In our case, the outer medium is the outer part of the progenitor star from which the magnetar formed (see §4); we do not neglect the thermal pressure, but we do assume that the progenitor is unmagnetized.

Given the evolution of H and p_n with time (calculated below), equations (8)-(13) were solved under the assumption of axisymmetry, to determine the evolution of the shape of

the shell with time. We were not able to cast these equations in full upwind form, because of the presence of a term describing the advection of the shell curvature, which is not constant and changes in time. This requires adding some artificial viscosity in order to avoid the numerical growth of perturbations.

One cautionary comment about the thin-shell model is in order. Equations (8)-(13) are momentum conserving, not energy conserving, in the sense that a shell expanding into an ambient medium has constant momentum and thus its energy decreases in time. The equations do conserve energy, however, in the sense that the work done by the interior bubble is self-consistently supplied to the shell (see § 4), but some of this energy is then lost as the shell expands and sweeps out into the ambient medium. Unfortunately, it is not possible to conserve both momentum and energy in the time dependent, thin-shell approximation (by contrast, in the self similar case discussed in the Appendix, one can satisfy both requirements, but in this case the time evolution is factored out of the equations). One consequence of this is that the calculations that follow probably evolve somewhat more slowly than would a true SN shock expanding into its host star, although we are confident that our conclusions about generating asymmetric bubbles are robust (e.g., the self-similar solutions in the Appendix show similar asymmetry).

4 A MAGNETAR IN A BOTTLE

In this section we use the thin-shell model to calculate the evolution of the magnetized bubble inflated by a central magnetar. As was pointed out in §2.1, one needs to derive the internal pressure distribution in the bubble in order to solve for the dynamics of the shell. In particular, one needs to know the value p_n of the total pressure on the axis and the value of the scale height H of the pressure distribution. Once these two parameters are known it is possible to derive the pressure throughout the bubble, in particular its value at the inner edge of the shell. One can show that given the shape of the shell bounding the bubble, p_n , and H , the total energy E_{tot} , the magnetic energy E_{mag} , and the magnetic flux Φ inside the bubble itself are uniquely defined, where

$$E_{tot} = \int_V p_n \mathcal{F}(r/H, z) dv, \quad (14)$$

$$E_{mag} = \int_V p_n \mathcal{G}(r/H, z) dv, \quad (15)$$

$$\Phi = \int_A \sqrt{p_n} \mathcal{Q}(r/H, z) da, \quad (16)$$

and where V is the volume of the bubble and A is the area in the $r - z$ plane, delimited by the shell. The dimensionless functions $\mathcal{F}, \mathcal{G}, \mathcal{Q}$ are given in terms of cylindrical coordinates, and can be derived from the pressure and magnetic field given by equations (5)-(6) in § 2.1.

In order to compute the evolution of the internal structure in the bubble we subdivided each time step (dt) of the shell evolution into two sub-steps. In the first sub-step, that we call adiabatic, we neglect injection of energy and magnetic field by the central source, and we compute the adiabatic losses due to expansion according to:

$$dE_{tot} = \int_S p dV, \quad (17)$$

where p is the total pressure along the shell surface S and dV is the volume increment that results from the evolution of the shell surface. Once the adiabatic losses are known one can derive the new value for the total energy in the bubble. During this adiabatic step the magnetic flux remains constant. After the adiabatic step, the new values of p_n and H are re-evaluated by solving the following equations:

$$E_{tot,a} = E_{tot} - dE_{tot} = \int_V p_n \mathcal{F}(r/H, z) dv, \quad (18)$$

$$\Phi = \int_A \sqrt{p_n} \mathcal{Q}(r/H, z) da, \quad (19)$$

where the integrals are computed using the values of V and A after the expansion. Using the new values of p_n and H , we need to recompute the new magnetic energy inside the bubble $E_{mag,a}$, because adiabatic losses act on the total energy. This is done using equation (15).

In the second sub-step, that we call the injection step, the shape of the bubble is assumed to be fixed and we compute the new values of the total energy and the magnetic energy given the rate of total energy and magnetic energy injection by the central magnetar. The two equations to be solved for p_n and H are:

$$E_{tot,a} + \dot{E}_{tot} dt = \int_V p_n \mathcal{F}(r/H, z) dv, \quad (20)$$

$$E_{mag,a} + \dot{E}_{mag} dt = \int_V p_n \mathcal{G}(r/H, z) dv, \quad (21)$$

and once p_n and H are known we can also recompute the magnetic flux Φ , which will be needed in the next time step. With this method we determine the evolution of the pressure on the inner edge of the shell as a function of time given $\dot{E}_{tot}(t)$ and $\dot{E}_{mag}(t)$ (by, e.g., the results of Figure 1).

Based on modeling the spectra of supernovae associated with nearby GRBs, there are some indications that GRBs arise from very massive stars with ZAMS masses of $M \approx 40 M_\odot$ (e.g., Iwamoto et al. 1998). There are also observational indications that Galactic magnetars are formed from comparably massive stars (Gaensler et al. 2005, Muno et al. 2006). We thus consider the evolution of a magnetized bubble inside a progenitor star of $35 M_\odot$, using the progenitor models of Woosley et al. (2002). We have also considered lower progenitor masses down to $\approx 11 M_\odot$, which may be more appropriate for the progenitors of X-ray flashes (Mazzali et al. 2006). We find little difference in the results for different progenitors, at the level of quantitative detail to which we aspire. The most significant effect is that for fixed energy injection, the bubble expands more slowly for more massive progenitors. The relative insensitivity to progenitor mass can in part be understood by noting that the self-similar solutions described in the Appendix show explicitly that the elongation of the bubble depends only weakly on the density profile of the ambient medium.

As discussed in §2, our model of the magnetar wind assumes that it is expanding into a cavity evacuated by the outgoing SN shock. To initialize our simulations, we thus carve out a spherical cavity with a radius of 10^9 cm inside our progenitor, corresponding to the region of infall in the first ~ 1 sec. We assume that this cavity is bounded by a

thin shell whose mass is equal to the mass that originally was in the cavity region minus $1.4 M_\odot$ (the canonical mass for a neutron star). In all of our simulations, time is defined after core bounce and the simulation starts 1 second after core bounce. Moreover we impart to the shell an outward velocity so that the total shell energy at the beginning is 10^{51} ergs, enough to trigger a SN. If instead one assumes an initially stationary shell, the evolution is essentially unchanged for weakly magnetized bubbles because the pressure of the bubble is relatively isotropic (this assumes that the magnetar wind extracts at least $\sim 10^{51}$ ergs at early times, as is the case in the models shown in Figure 1). For strong magnetization, the elongation of the bubble along the axis is also nearly independent of the initial shell energy. However, for large E_{mag}/E_{tot} , the pressure in the bubble near the equator can be so small that infall cannot be prevented. To model this case, a full hydrodynamic solution is required.

We follow the evolution of the shell and interior bubble to large distances, into the hydrogen envelope of the progenitor. For GRB progenitors, the hot plasma confined inside will emerge into the circumstellar medium once the shell surface reaches the outer edge of the helium core. The initial material that emerges will probably only move with modest Lorentz factor. Subsequent material will, however, rapidly accelerate through the channel carved by the magnetar wind, reaching asymptotic Lorentz factors set roughly by the enthalpy of the material in the bubble (assuming that $E_{mag} \lesssim E_{thermal}$ in the bubble). This phase of evolution cannot be studied using the thin shell-approximation, but requires full relativistic MHD simulations. Nonetheless, it appears natural that a highly relativistic and collimated outflow will emerge out of the cavity carved by the early magnetized bubble.

4.1 Results

In Figure 4 we show the results of a series of simulations for different fixed values of $\dot{E}_{mag}/\dot{E}_{tot}$, the ratio of the Poynting flux to the total power injected by the magnetar. In all of the calculations in Figure 4, we assume that the total power supplied by the central source is given by $\dot{E}_{tot} = 10^{51} (t/1 \text{ s})^{-1} \text{ erg s}^{-1}$, which is a reasonable approximation to the lower power solutions in Figure 1 (note that we neglect the possibility of early injection of energy and start our simulation 1 second after core bounce). Note that in this case equal energy is supplied per decade in time.

For values of $\dot{E}_{mag}/\dot{E}_{tot} \sim 0.1$, the pressure distribution inside the bubble is relatively spherical (Figs. 2 & 3) and so the surrounding shell becomes only modestly asymmetric. Most of the energy supplied by the central magnetar in this case is transferred to the surrounding SN shock and host star. Low magnetization bubbles of this kind would thus likely produce a mildly asymmetric hyper-energetic SNe, but it appears unlikely that the relativistic material supplied by the magnetar can easily escape its host star. For larger values of $\dot{E}_{mag}/\dot{E}_{tot}$, the shell evolves more asymmetrically because most of the pressure is exerted along the axis for magnetized bubbles. By $\dot{E}_{mag}/\dot{E}_{tot} \gtrsim 0.3$, there is clear evidence for a very elongated channel driven through the host star by the anisotropic pressure of the central bubble. The shell reaches the outer edge of the progenitor ($\sim 2 \cdot 10^{10}$ cm) after $\approx 5 - 10$ sec. At this point the ambient density drops to typical




Figure 4. Evolution of a magnetized bubble inside a $35M_{\odot}$ progenitor for a central source with $\dot{E}_{tot} = 10^{51} (t/1s)^{-1} \text{ erg s}^{-1}$. The initial time is 1s; the contours describe the shape of the shell at 1s time intervals for the first three panels and 0.5s time intervals for the last. From left to right, the solutions correspond to increasing the magnetization of the bubble, with $\dot{E}_{mag}/\dot{E}_{tot} = 0.1, 0.2, 0.3$, & 0.4 , respectively. For $\dot{E}_{mag}/\dot{E}_{tot} \gtrsim 0.3$ most of the pressure of the inner bubble is exerted close to the axis (see Figs. 2 & 3), which leads to the asymmetric evolution of the bounding shell.

circumstellar values, and the shell will rapidly blow out of the star. The highly relativistic material contained in the interior bubble can now flow relatively unimpeded out of the host star, forming a relativistic jet; it is natural to associate these models with the production of a GRB.

Figure 5 shows the evolution of the thin shell for a more energetic, but more rapidly decaying, central source with $\dot{E}_{tot} = 10^{52} (t/1s)^{-2} \text{ erg s}^{-1}$, which is an approximation to the higher power solutions in Figure 1. We consider $\dot{E}_{mag}/\dot{E}_{tot} = 0.2$ (left) and $\dot{E}_{mag}/\dot{E}_{tot} = 0.3$ (right). Note that in this case, most of the energy is supplied to the bubble at early times and so the evolution of the system is similar to the case of a magnetic bomb with a fixed energy of $\sim 10^{52}$ ergs in the bubble. The evolution of the shell in Figure 5 is qualitatively similar to that of the lower power solutions shown in Figure 4, although the bubble evolves more rapidly because of the more energetic central source. One consequence of this more rapid evolution is that the shell velocity is closer to c , implying that the assumption of magneto-hydrostatic equilibrium used to derive the interior structure is likely to be less accurate than in the case of the weaker power sources in Figure 4.

For PNSs with rotation periods longer than the values of $\approx 1 - 2$ ms considered in Figure 1, the energy injection rate will be lower and approximately constant at early times because the spindown time is longer than the Kelvin-Helmholtz time of $\approx 10 - 100$ s. To investigate this limit, we considered the evolution of a bubble with a constant energy injection rate of $\dot{E}_{tot} \approx 10^{50} \text{ erg/s}$. Elongation analogous to that shown in Figures 4 & 5 can be achieved, although somewhat higher magnetization is required. An asymmetry similar to the $\dot{E}_{mag}/\dot{E}_{tot} = 0.2$ solution in Figure 4 requires $\dot{E}_{mag}/\dot{E}_{tot} = 0.3$ for this lower $\dot{E}_{tot}$ and takes a somewhat longer time ~ 20 sec to develop. This example highlights that lower power sources – which can originate from more modestly rotating PNSs – can still lead to asymmetric bubbles because the energy per unit solid angle along the pole is significant even for modest $\dot{E}_{tot} \sim 10^{49} - 10^{50} \text{ erg s}^{-1}$. Such




Figure 5. Evolution of a magnetized bubble inside a $35M_{\odot}$ progenitor for a central source with a total spindown power of $10^{52} (t/1s)^{-2} \text{ erg s}^{-1}$; the initial time is 1s. Contours represent the shape of the shell at 1s intervals. *Left:* $\dot{E}_{mag}/\dot{E}_{tot} = 0.2$ *Right:* $\dot{E}_{mag}/\dot{E}_{tot} = 0.3$.

sources may form asymmetric SN and, in some cases, very long-duration GRBs or X-ray flashes.

An approximate analytic understanding of the late-time structure of the shell shown in Figures 4 & 5 can be obtained by assuming that most of the energy is released inside the bubble before it has time to expand significantly – so that its shape is still approximately spherical and adiabatic

losses can be neglected. In this case most of the pressure will reside in a region along the axis whose opening angle is $\approx H/R$ and the shell will expand rapidly along the axis within this solid angle. Figure 3 gives the relationship between H/R and the magnetization of the bubble, which can be used to estimate the opening angle of the resulting “jet” at late-times. For example, Figure 3 shows that $H/R \approx 0.1$ for $E_{\text{mag}}/E_{\text{tot}} \approx 0.3 - 0.4$, which is reasonably consistent with the angular collimation of ~ 0.1 rad in Figure 4. It is also worth noting that the high axial pressure produced by a magnetized bubble with energy E leads to an expansion along the axis that is quantitatively similar to that produced by a jet with kinetic power

$$L_{\text{jet}} \simeq 7.5 \cdot 10^{51} \theta \left(\frac{E}{10^{51} \text{erg}} \right) \left(\frac{10^9 \text{cm}}{R} \right) \text{erg s}^{-1}, \quad (22)$$

where we have assumed that the angular collimation $\theta \sim H/R$ and that the dimensionless axial pressure $\bar{P} \approx R/H$ (which is reasonable for $H \gtrsim 0.03 R$; Figure 3).

The results in Figures 4 & 5 assume that $\dot{E}_{\text{mag}}/\dot{E}_{\text{tot}}$ is independent of time. This may be a poor approximation given the strong evolution in the magnetization of the wind as a function of time at small radii (Fig. 1). Indeed, one might naively expect that $\dot{E}_{\text{mag}}/\dot{E}_{\text{tot}}$ would increase in time on a timescale of a few sec, given the evolution of $\sigma_{LC}(t)$ for magnetar winds. If this is correct, the magnetar-created bubble may initially impart its energy relatively spherically, enhancing the energy of the outgoing SN shock (as in the left panel of Figure 4). At late times, however, the bubble will drive a jet out along the axis (as in the right-most panel in Figure 4). To explore this scenario, we carried out a series of simulations starting with a small initial value of $\sigma \approx 0.05 - 0.1$ and then increasing σ in time as σ_{LC} increases, up to the equipartition value of $\sigma \approx 0.5$. As expected, in the first few seconds the evolution of the shell was quite spherical, closely resembling the $\sigma = 0.1$ panel of Fig. 4, while at late times the shell evolves into a more elongated structure analogous to the $\sigma = 0.3 - 0.4$ panels of Fig. 4. In this scenario, the different panels in Fig. 4 qualitatively describe different snapshots in time for the evolution of a shell driven by a wind with increasing magnetization. This suggests that the increasing magnetization of the magnetar wind provides a way of tapping the spindown energy to generate a mildly asymmetric hyper-energetic SN, while at the same time creating an axial cavity along which relativistic material can escape, powering a GRB.

Throughout this paper, we have considered only models in which the progenitor has a spherical density profile, in order to understand the importance of magnetic stresses on the asymmetric evolution of the stellar wind bubble. However for rotating stars the density in the polar region is expected to be smaller than at the equator. This can also facilitate collimation. A full investigation of the combined collimation induced by magnetic stresses and a non-spherically symmetric stellar profile is beyond the scope of this paper. We have, however, carried out a few preliminary calculations investigating the two extremes: a low density polar channel whose opening angle is either smaller or bigger than H/R_{polar} . Our results show that at low magnetization the presence of a low density channel along the rotation axis can lead to significant collimation (well in excess of the mild asphericity shown in

Figure 4 at low magnetization), while for higher magnetization the collimation is primarily due to the magnetic stresses we have focused on in this paper.

5 DISCUSSION

In this paper we have calculated the evolution of a magnetized bubble formed inside an exploding massive star. Our motivation is to understand the impact of a newly born millisecond magnetar on its surrounding stellar envelope, and in particular to determine the conditions under which magnetar birth can produce the collimated outflows inferred from observations of long-duration GRBs.

Neutron stars born with $P \sim 1$ ms and $B \sim 10^{15} - 10^{16}$ G lose $\sim 10^{51} - 10^{52}$ ergs in $\sim 10 - 100$ sec in a magnetized wind that becomes increasingly relativistic at late times (Figure 1). This energy forms a bubble of plasma and magnetic field confined by the inertia of the surrounding star. If the material supplied to the bubble has low magnetization, $\dot{E}_{\text{mag}}/\dot{E}_{\text{tot}} \lesssim 0.1$, the bubble expands relatively spherically (Figure 4) and most of the energy of the spinning down neutron star is transferred to the surrounding SN shock, plausibly powering a hyper-energetic SN, but probably not a GRB. By contrast, for more appreciable magnetization, $\dot{E}_{\text{mag}}/\dot{E}_{\text{tot}} \gtrsim 0.3$, the bubble created by the magnetar rapidly becomes highly elongated along the rotation axis of the system as a result of the internal pressure distribution (§2.1), forming a cavity out of which the late-time relativistic wind from the magnetar can escape (Figure 4 & 5). We suggest that this is plausible physical mechanism for forming collimated relativistic outflows from newly formed millisecond magnetars.³

This mechanism works even if the host star is spherically symmetric. In addition, even if most of the wind energy flux is concentrated in the equatorial plane (as is expected for relativistic winds from neutron stars), the surrounding bubble will still reach magnetohydrostatic equilibrium and will thus elongate along the axis due to magnetic stresses as we have described. Finally, we note that it is not necessary to wait until late times, when the magnetar wind is relativistic, in order for the bubble to evolve highly asymmetrically. Collimation can be achieved in the early mass loaded phase, provided only that the bubble contains a sufficient toroidal magnetic field. This mechanism may then operate in addition to hoop-stress collimation of the non-relativistic flow (e.g., Bucciantini et al. 2006; Uzdensky & MacFadyen 2006). This early time non-relativistic phase cannot by itself produce a GRB, but can create a channel out of which the later relativistic wind emerges. Such a channel might also provide boundary conditions conducive to the acceleration of the wind and the conversion of magnetic energy

³ Although we have framed much of this paper in terms of the magnetar model for GRBs, magnetized outflows from an accretion disk around a neutron star or black hole would produce asymmetric magnetized bubbles akin to those considered in this paper. Whether they would play a similar role in driving collimated flows inside a supernova depends on the details of the disks’ mass and magnetic energy losses as a function of time, as well as uncertainties in the disk winds’ σ similar to those encountered in the neutron star winds considered here.

into kinetic energy (Komissarov et al. 2007). Our calculations show that for the expected magnetar energy loss rates, a collimated cavity is formed after ~ 10 sec (Fig. 4). At this point, magnetar winds have $\sigma_{LC} \sim 100$ (Fig. 1), in the range required to account for GRBs.

Because the birth rate of magnetars ($\sim 10\%$ of neutron stars; e.g., Kouveliotou et al. 1994) is significantly larger than the rate of GRBs ($\sim 0.1 - 1\%$ of massive stellar deaths; e.g., Podsiadlowski et al. 2004), most magnetar births cannot produce standard long-duration GRBs. This is presumably either because an extended stellar envelope inhibits the escape of a collimated outflow or because most magnetars are born rotating more slowly than the millisecond rotators we have focused on in this paper. For more modestly rotating PNSs, the asymmetric expansion of a magnetized bubble could contribute to the inferred asymmetry of many core-collapse supernovae (e.g., Wang et al. 2001). In addition, a PNS with, e.g., $P \approx 4$ ms and $B \approx 3 \times 10^{15}$ G has a rotational energy of $\approx 10^{51}$ ergs and a spindown time of ≈ 1 day. The birth of such a neutron star would not produce a hyper-energetic SN or a canonical GRB. However, if the bubble created by the magnetar is sufficiently magnetized, it would evolve asymmetrically in a manner similar to the calculations shown in Figures 4 & 5. This could produce a long-duration transient analogous to the X-ray flash 060218 associated with SN 2006aj (Mazzali et al. 2006; Soderberg et al. 2006; we should note, however, that many X-ray flashes may have lower inferred energies because of viewing angle effects rather than being intrinsically less energetic events; e.g., (Granot et al. 2005). The remnant Cass A, with its strong jet/counter-jet morphology (e.g., Hwang et al. 2004), may be an example of an asymmetric explosion driven and shaped by a magnetized wind accompanying magnetar birth. Indeed, Chakrabarty et al. (2001) suggested that the central X-ray point source in Cass A is a magnetar.

The thin-shell calculations described in this paper assume that the magnetar wind expands into an initially spherical cavity created by an outgoing SN shock. This requires that the spindown time of the magnetar is at least somewhat longer than the time required to initiate the stellar explosion (i.e., $\gtrsim 1 - 2$ sec). Our assumption of a “successful” SN explosion does not, of course, preclude that the explosion itself is magneto-centrifugally driven, as in the force-free model for the collimated explosion of a star by a newly-formed magnetar in an otherwise “failed” SN (e.g., Ostriker & Gunn 1971 or Uzdensky & MacFadyen 2007). However, one interesting problem not addressed by our calculations is the spindown of the magnetar and the evolution of its surrounding bubble if the initial explosion is primarily bipolar (see, e.g., the simulations of Moiseenko et al. 2006 and Burrows et al. 2007). Late-time collimation of relativistic material in this context may be modified by the large inertia of the accreting stellar envelope (or fallback material) in the equator of the star (see also the related arguments of Uzdensky & MacFadyen 2007). In addition, it is worth noting that if the outflow always has high magnetization, our calculations suggest that, because most of the pressure will be exerted along the axis, there could be a collimated GRB but no associated equatorial explosion. This could account for the recently discovered supernova-less GRBs (Fynbo et al. 2006).

One of the uncertainties associated with our calculations is that the magnetization of the material supplied to the surrounding bubble is difficult to calculate. Magnetic energy has to be supplied to the bubble relatively rapidly, with $\sigma = \dot{E}_{mag}/\dot{E}_{tot} \gtrsim 0.2$ at the termination shock in our models that show significant collimation. Observations of PWNe suggest quite low $\sigma \sim 0.01$ at the termination shock, which would imply that there is insufficient time to build up the anisotropic magnetic stress needed to drive aspherical expansion of the surrounding stellar envelope. However, we suspect that the confined bubbles around newly formed magnetars will have higher magnetization at their termination shocks than has been inferred in PWNe and in pulsar-Be star binaries. This is because the distance to the termination shock is only ~ 10 light cylinder radii in our problem, relative to $> 10^4$ light cylinder radii in the systems where we have direct observational constraints. As a result, there is less time for the magnetic field in the wind to dissipate, plausibly leading to higher magnetization.

All of the calculations described in this paper are based on the thin-shell approximation. This model is useful for demonstrating that magnetar birth can produce conditions conducive to the formation of a collimated outflow that can emerge out of the host star. However, modeling this process in detail is beyond the scope of the present simplified calculations, and will require full relativistic MHD simulations. Indeed, it is our intention to use the results of the present paper as a guide for more realistic simulations. Such calculations are necessary to determine the fraction of the spindown energy that goes into a relatively spherical explosion of the host star relative to the energy that flows out of the collimated cavity. Quantifying this is important for understanding the conditions under which magnetar birth might produce *both* a hyper-energetic SN and a GRB, as is observed (e.g., Woosley & Bloom 2006). We have speculated in §4 that this can occur if the surrounding bubble becomes progressively more magnetized as the magnetar spins down, but multi-dimensional simulations are needed to assess this (and to understand if it is a necessary condition). Multi-dimensional simulations will also allow a more detailed study of how the late-time relativistic outflow emerges from within the earlier non-relativistic wind and the different observational signatures associated with each phase of the outflow (analogous to studies of jets emerging from a host star; e.g., Morsony et al. 2007). They also allow investigation of the stability of the confining envelope, which is subject to possible Rayleigh-Taylor fragmentation, since it is accelerated by the light weight bubble (e.g., Arons 2003), an effect likely to be of substantial significance at higher energy injection rates. Such instabilities could be important for understanding the short timescale variability observed in GRBs, as could intrinsic time variability in the magnetar wind, driven by, e.g., reconnection in the equatorial current sheet (Bucciantini et al. 2006). Variability might also originate inside the bubble due to the dynamics of the interaction of the wind with the progenitor (as is seen in PWNe simulations; e.g., Del Zanna et al. 2004). Finally, in addition to the asymmetric expansion of the entire bubble discussed here, observations and simulations of PWNe also reveal a moderately relativistic axial “jet” within the nebula itself (e.g., Weisskopf et al. 2000; Del Zanna et al. 2004); this may be dynamically important for proto-magnetar bubbles as well.

ACKNOWLEDGMENTS

N.B. was supported by NASA through Hubble Fellowship grant HST-HF-01193.01-A, awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS 5-26555. EQ and BDM were supported by the David and Lucile Packard Foundation and a NASA GSRP Fellowship to BDM. JA's research has been supported by NSF grant AST-0507813, NASA grant NNG06GI08G, and DOE grant DE-FC02-06ER41453, all at UC Berkeley; by the Department of Energy contract to the Stanford Linear Accelerator Center no. DE-AC3-76SF00515; and by the taxpayers of California. We thank A. Heger for making massive stellar progenitor models available online.

APPENDIX A: SELF-SIMILAR SOLUTIONS

It is well known that the 1D thin shell equations admit self-similar solutions, where the time dependence can be factored out. Chevalier & Luo (1994) have shown that equations (8)-(13) can be solved in the simplified case of a wind blowing inside a wind, when the shell expands at a constant speed. In this case it is possible to reduce equations (8)-(13) to a system of ordinary differential equations. Here we extend their treatment by considering the expansion of a hot magnetized bubble inside either a stationary medium (as in the case of the ISM or the roughly static envelope of a progenitor star) or freely expanding ejecta (as in the case of PWNe). We derive the conditions under which self-similarity applies and discuss how the elongation of the bubble scales with its magnetic energy content. We limit our investigation to the case of an isotropic outer medium, even though self-similarity holds in the more general case of fixed latitudinal dependence (Kahn & West 1985, Campbell et al. 2003).

If one assumes a cold unmagnetized outer medium with a power law density profile $\rho_o \propto r^\alpha$, that $p_n \propto t^\chi$ and that R_{polar}/H is constant, where R_{polar} is the radius of the shell along the axis, one can search for a solution of the form

$$\begin{aligned} R &= R(\theta)t^\beta \\ v_\perp &= v_\perp(\theta)t^\delta \\ v_\parallel &= v_\parallel(\theta)t^\delta \\ \sigma &= \sigma(\theta)t^\eta \end{aligned}$$

Taking $\delta = \beta - 1$, $\eta = \beta + \alpha\beta$, and $\chi = 2\beta + \alpha\beta - 2$, the time dependence can be eliminated and the problem reduces to a system of ordinary differential equation in the θ direction. Neglecting the outer pressure and magnetic field, and the inner ram pressure, and assuming a stationary outer medium (a similar derivation can be obtained for freely expanding ejecta), equations (8)-(13) can be written as

$$\tan \xi = -\frac{1}{R} \frac{\partial R}{\partial \theta}, \quad (A1)$$

$$\beta\sigma = \rho_o(\beta R \cos \xi) - 2\beta\sigma -$$

$$\frac{\cos \xi}{\sin \theta R^2 \partial \theta} [R \sin \theta \sigma (v_\parallel - \beta R \sin \xi)] \quad (A2)$$

$$\beta(\beta - 1)\sigma R \cos \xi = -\rho_o \beta^2 R^2 \cos^2 \xi + (p_i + B_i^2/8\pi) -$$

$$\sigma[v_\parallel + \beta R \sin \xi] \frac{\cos \xi}{R} \left[\frac{\partial}{\partial \theta} (\beta R \cos \xi) - v_\parallel \left(1 + \frac{\partial \xi}{\partial \theta} \right) \right] \quad (A3)$$

$$(\beta - 1)\sigma v_\parallel = -\rho_o \beta R \cos \xi v_\parallel -$$

$$\sigma[v_\parallel + \beta R \sin \xi] \frac{\cos \xi}{R} \left[\frac{\partial}{\partial \theta} (v_\parallel) + \beta R \cos \xi \left(1 + \frac{\partial \xi}{\partial \theta} \right) \right], \quad (A4)$$

where the variables are now only function of θ , and $p_i + B_i^2/8\pi = p_n f(R \sin \theta/H)$. Note that these equations are not easy to write in non-dimensional form because, as we will discuss below, the characteristic length-scale is an eigenvalue of the problem. Only in the case $\alpha = -2$ is the solution relatively straightforward (Chevalier & Luo 1994).

To check the validity of the self-similar solution one must verify a posteriori that R_{polar}/H does not change in time and that the pressure p_n scales as a power law. We first consider the scale height H . As discussed in §4, the evolution of the bubble can be described as a series of infinitesimal adiabatic expansions followed by energy injection at fixed volume. It is easy to show that, in the relativistic case, when the adiabatic index is 4/3, entropy and magnetic flux conservation imply that the ratio R_{polar}/H is indeed constant during self-similar expansion. On the other hand, as long as the ratio of the Poynting flux to the total luminosity, $\dot{E}_{mag}/\dot{E}_{tot}$, is equal to the ratio of the magnetic energy to total energy in the bubble E_{mag}/E_{tot} , the height scale H is constant, at fixed shape.

Despite the fact that the thin-shell equations conserve only momentum, it is possible to conserve energy as well by requiring the correct temporal scaling, in which case the total energy in the system increases only because of injection ($\dot{E}_{tot}$). The total energy in the shell is:

$$\int_S \frac{1}{2} \sigma (v_\parallel^2 + v_\perp^2) dS = E_{sh} t^{5\beta + \alpha\beta - 2}, \quad (A5)$$

where S is the shell surface and E_{sh} is a constant that depends on the shell structure. If the ratio R_{polar}/H is constant then the total energy in the bubble is

$$\int_V e(r, z) dv = p_n K t^{3\beta}, \quad (A6)$$

where V is the volume of the bubble, K is a constant depending on the shell shape and the ratio R_{polar}/H , and e is the energy density in the bubble. If $p_n \propto t^\chi$ then the total internal energy scales as the shell energy. Thus one can write the total energy in the system as

$$E_{tot}(t) = \int_o^t \dot{E}_{tot} dt \propto t^{5\beta + \alpha\beta - 2}. \quad (A7)$$

This relation shows that as long as $\dot{E}_{tot}$ is a power-law in time, self similar solutions are self-consistent. Equation (A7) together with the ratio $\dot{E}_{mag}/\dot{E}_{tot}$ determines how the self-similar shape is related to the injection properties, namely the total luminosity and Poynting flux. Once α and R_{polar}/H are given one can search for a solution of equations (A1)-(A4).

A0.1 Numerical Methods

In the case studied by Chevalier & Luo (1994), the shell moves at a constant speed, $\beta = 1$, and the system of ordinary differential equations can be easily solved as an initial value problem, assuming regular conditions on the axis. Given that the equations are singular on the axis, it is necessary to expand them in Taylor series and solve for the

coefficients. In more general cases, the shell is subject to acceleration and $\beta \neq 1$. This changes the nature of the equations and makes the problem much harder to solve. If one tries to expand the equations in a Taylor series on the axis, which works for the case $\beta = 1$, one finds that terms of order $\mathcal{O}(\theta^n)$ depend on terms of order $\mathcal{O}(\theta^{n+1})$, and the series is open. This is typical of two-point boundary value problems (eigenvalue problems) where conditions at one boundary are not sufficient. However it is not obvious where and what kind of additional boundary conditions one should impose. One might enforce regularity on the equator, either by assuming a smooth shape or zero parallel velocity. However the solutions of Chevalier & Luo (1994) show that discontinuities at the equator might arise, especially for larger values of R_{polar}/H . We have chosen to impose that the tangential transport velocity should be 0 on the equator to avoid the formation of an equatorial ring where matter can accumulate indefinitely, but this does not preclude other possible boundary conditions.

We tried several methods for solving the steady state angular equations given by (A1)-(A4). These included standard shooting techniques, using both explicit ODE solvers and an implicit integrator, standard relaxation methods, and a Fourier expansion of the angular equations. For a variety of reasons, we find that none of these methods was satisfactory for finding solutions, aside from the relatively spherical limit when $R_{\text{polar}}/H \lesssim 1$. Instead, we found that the most successful method was to use the time dependent equations (eqs. [8]-[13]) to evolve the system forward in time, imposing a fixed value for R_{polar}/H and a pressure $p_n \propto t^\chi$, where the exponent χ is the one expected for the self-similar solution. Self-similarity is then reached at late times after the initial transient dies away; we have verified that the shell at late times has converged to a self-similar shape, with only a few percent error in radius. It is, however, known that accelerated shells are subject to corrugational instabilities (the thin-shell instability). We have included an artificial viscosity to suppress this instability and focus on the overall evolution of the bubble. Instabilities may indeed be important for the evolution of the shell but we leave a study of them to future work. In each case, we have determined a posteriori that the chosen value of viscosity was small enough not to affect the shape of the bubble. This method for determining the self-similar solution worked up to $R_{\text{polar}}/H \sim 10$. More elongated bubbles were difficult to investigate unless one started with initial conditions very close to the desired self-similar solution. Otherwise the deviation from the correct self-similar solution was effectively a large-amplitude large-scale initial perturbation; in this case, we found that the corrugational instability could not be suppressed without increasing the viscosity to a point where it modified the overall shape of the bubble.

A0.2 Results

In Figure A1 we plot the shape of the shell for three different values of R_{polar}/H , and for $\alpha = 0, -1, -2$ in the case of a constant injection luminosity ($\beta = 3/(5+\alpha)$). For $\alpha = 0$ and -1 , the solutions are found by the methods described in the previous subsection and are restricted to $R_{\text{polar}}/H \lesssim 10$. The $\alpha = -2$ case is that of Chevalier & Luo (1994) for which the angular equations (A1)-(A4) can be straightfor-

wardly integrated. In this case, we compute solutions up to $R_{\text{polar}}/H \approx 100$, for comparison to the time-dependent solutions in the main text.

Figure A1 shows that the elongation of the bubble increases with increasing R_{polar}/H , and for smaller values of α . In the case of a constant density outer medium, the shell appears to be regular both at the pole and at the equator, while in the case $\alpha = -1$ and -2 for large values of R_{polar}/H , a cusp is formed in the equatorial plane, where matter tends to accumulate. We have verified that all quantities scale according to self similarity.

Unlike in the case studied by Chevalier & Luo (1994), in which the surface density monotonically increases from the pole to the equator, for $\alpha = 0$ and small values of R_{polar}/H we find that the surface density reaches a maximum at intermediate latitudes.

Figure A2 shows a summary of the results of our simulations for the case of constant injection power into a static outer medium (typical for wind bubbles): shown are the ratio of the magnetic energy to the total energy and the elongation of the bubble, as functions of R_{polar}/H . The elongation scales approximately as $\sqrt{R_{\text{polar}}/H}$ and, as expected, is larger for steeper density profiles. The dependence on the density profile of the outer medium is, however, quite weak, with less than a factor of 2 difference between $\alpha = 0$ and $\alpha = -2$. For the magnetar problem considered in this paper, this suggests that the outer density structure of the progenitor star should not significantly effect the collimation of the bubble (this is consistent with the fact that the asymmetry of the bubble was only a weak function of the progenitor star, as discussed in §4). Figure A2 also shows that in all cases the ratio $E_{\text{mag}}/E_{\text{tot}}$ inside the bubble is relatively small, and tends to be smaller for smaller values of α . This demonstrates that strong elongation does not require magnetically dominated bubbles, which is consistent with our result for the evolution of magnetized bubbles inside GRB progenitors in §4.

We also carried out simulations for a bubble inside freely-expanding ejecta with velocity $v_o \propto R/t$ and density $\rho \propto r^\alpha t^{-\alpha-3}$. We find that for $\alpha = -2$, the time-dependent solution never reaches a self-similar solution, but instead the evolution always depends on the initial conditions; for this reason, we do not show solutions for $\alpha = -2$. This is because in the case $\alpha = -2$ if one assumes the expected power law temporal evolution for the internal pressure, it is easy to show that the ratio between the inner pressure and the ram pressure of the outer medium is constant, and does not depend on the radial evolution. In this sense, the bubble never relaxes to self-similarity. The case of freely-expanding ejecta is relevant for a central source which is not energetic enough to produce a bubble which effects the dynamics of the stellar explosion on short time-scales, but instead creates an energetic PWN inside the expanding SN ejecta. This could be relevant to young PWN generally and also to lower energy transients such as X-ray flashes (see §5). The results for the elongation of the bubble and its dependence on the magnetization are shown in Figure A3; they are reasonably similar to the results shown in Figure A2.




Figure A1. Self-similar shapes for magnetically driven bubbles expanding into a static ambient medium with $\rho \propto r^\alpha$. Results are shown for three different values of R_{polar}/H . Results for $\alpha = -2$ are based on the model of Chevalier & Luo (1994) while for $\alpha = 0$ and -1 , they are based on time dependent thin-shell simulations that reach self-similarity at late times.

REFERENCES

- Arons, J. 2003, *ApJ*, 589, 871
- Begelman, M. C., & Li, Z. 1992, *ApJ*, 397, 187
- Berger, E., Kulkarni, S. R., & Frail, D. A. 2004, *ApJ*, 612, 966
- Blasi, P., Epstein, R.I., and Olinto, A.V. 2000, *ApJ*, 533, L123
- Bucciantini, N. 2002, *A&A*, 387, 1066
- Bucciantini, N., Thompson, T. A., Arons, J., Quataert, E., & Del Zanna, L. 2006, *MNRAS*, 368, 1717
- Bucciantini, N., Thompson, T. A., Arons, J., Quataert, E., & Del Zanna, L. 2007, *Proceedings of the 2006 COSPAR Meeting in Beijing*, accepted for publication in *AdSR*, *ArXiv Astrophysics e-prints*, arXiv:astro-ph/0701578
- Burrows, A. 1986, *ApJ*, 300, 488
- Burrows, D. N., & Racusin, J. 2007, *ArXiv Astrophysics e-prints*, arXiv:astro-ph/0702633
- Burrows, A., Dessart, L., Livne, E., Ott, C. D., & Murphy, J. 2007, *ArXiv Astrophysics e-prints*, arXiv:astro-ph/0702539
- Campbell, J. R., Borkowski, K. J., & Blondin, J. 2003, *Bulletin of the American Astronomical Society*, 35, 613
- Canto, J. 1980, *A&A*, 86, 327
- Canto, J., Raga, A. C., & Wilkin, F. P. 1996, *ApJ*, 469, 729
- Chakrabarty, D., Pivovarov, M. J., Hernquist, L. E., Heyl, J. S., & Narayan, R. 2001, *ApJ*, 548, 800
- Chevalier, R. A., & Luo, D. 1994, *ApJ*, 421, 225
- Coroniti, F. V. 1990, *ApJ*, 349, 538
- Del Zanna, L., Amato, E., & Bucciantini, N. 2004, *A&A*, 421, 1063
- Duncan, R. C., Shapiro, S. L., & Wasserman, I. 1986, *ApJ*, 309, 141
- Duncan, R. C., & Thompson, C. 1992, *ApJL* 392, L9
- Eichler, D., & Waxman, E. 2005, *ApJ*, 627, 861
- Frail, D. A., et al. 2001, *ApJL*, 562, L55
- Fynbo, J. P. U., et al. 2006, *Nature*, 444, 1047
- Gaensler, B.M., Arons, J., Kaspi, V.M., et al. 2002, *ApJ*, 569, 878
- Gaensler, B. M., McClure-Griffiths, N. M., Oey, M. S., Haverkorn, M., Dickey, J. M., & Green, A. J. 2005, *ApJL*, 620, L95
- Giuliani, J. L. 1982, *ApJ*, 256, 624
- Goldreich, P., & Julian, W. H. 1970, *ApJ*, 160, 971
- Granot, J., Ramirez-Ruiz, E., & Perna, R. 2005, *ApJ*, 630, 1003
- Heger, A., Woosley, S. E., & Spruit, H. C. 2005, *ApJ*, 626, 350
- Hwang, U., et al. 2004, *ApJL*, 615, L117
- Iwamoto, K., et al. 1998, *Nature*, 395, 672




Figure A2. Properties of magnetized bubbles expanding into a stationary medium with a power law density profile with $\rho \propto r^\alpha$, for $\alpha = 0$ (squares), $\alpha = -1$ (triangles), and $\alpha = -2$ (diamonds). The $\alpha = -2$ results are based on solving the steady state angular equations (A1)-(A4), which can be solved for larger values of R_{polar}/H than the $\alpha = 0$ and -1 cases, which are based on solving the time dependent thin shell equations. $E_{\text{mag}}/E_{\text{tot}}$ is the ratio between the magnetic and total energy in the bubble. The elongation is defined as the ratio between the length of the bubble along the axis and the maximum extent of the bubble away from the axis.




Figure A3. Same as Figure A2 but for the case of expansion into the freely expanding ejecta of a supernova remnant: $v_o \propto R/t$, $\rho \propto r^\alpha t^{-\alpha-3}$; results for $\alpha = 0$ (squares) and $\alpha = -1$ (triangles) are shown.

Kahn, F. D., & West, K. A. 1985, MNRAS, 212, 837

Kennel, C. F., & Coroniti, F. V. 1984, ApJ, 283, 694

Kirk, J. G., & Skjæraasen, O. 2003, ApJ, 591, 366

Komissarov, S. S., & Lyubarsky, Y. E. 2004, MNRAS, 349, 779

Komissarov, S. S., Barkov, M. V., Vlahakis, N., Konigl, & A. L. 2007, ArXiv Astrophysics e-prints, arXiv:astro-ph/0703146

Kouveliotou, C., et al. 1994, Nature, 368, 125

Lamers, H. J. G. L. M., & Cassinelli, J. P. 1999, Introduction to Stellar Winds, by Henny J. G. L. M. Lamers and Joseph P. Cassinelli, pp. 452. ISBN 0521593980. Cambridge, UK: Cambridge University Press, June 1999.

LeBlanc, J. M., & Wilson, J. R. 1970, ApJ, 161, 541

Lyubarsky, Y. E., & Eichler, D. 2001, ApJ, 562, 494

Lyubarsky, Y. E., & Kirk, J. G. 2001, ApJ, 547, 437

MacFadyen, A. I., & Woosley, S. E. 1999, ApJ, 524, 262

Mazzali, P. A., et al. 2006b, Nature, 442, 1018

Mazzali, P. A., et al. 2006a, ApJ, 645, 1323

Metzger, B. D., Thompson, T. A., & Quataert, E. 2007, ApJ, 659, 561

Michel, F. C. 1969, ApJ, 158, 727

Moiseenko, S. G., Bisnovaty-Kogan, G. S., & Ardeljan, N. V. 2006, MNRAS, 370, 501

Morsony, B. J., Lazzati, D., & Begelman, M. C., 2007, ApJ submitted (astro-ph/0609254)

Muno, M. P., et al. 2006, ApJL, 636, L41

Ostriker, J. P., & Gunn, J. E. 1971, ApJL, 164, L95

Pavlov, G. G., Kargaltsev, O. Y., Sanwal, D., & Garmire, G. P. 2001, ApJ, 554, L189

Podsiadlowski, P., Mazzali, P. A., Nomoto, K., Lazzati, D., & Cappellaro, E. 2004, ApJL, 607, L17

Pons, J. A., Reddy, S., Prakash, M., Lattimer, J. M., & Miralles, J. A. 1999, ApJ, 513, 780

Qian, Y.-Z., & Woosley, S. E. 1996, ApJ, 471, 331

Sato, G., et al. 2007, ApJ, 657, 359

Soderberg, A. M., et al. 2006, Nature, 442, 1014

Stevens, I. R., Blondin, J. M., & Pollock, A. M. T. 1992, ApJ, 386, 265

Thompson, C., & Duncan, R. C. 1993, ApJ, 408, 194

Thompson, C. 1994, MNRAS, 270, 480

Thompson, T. A., Burrows, A., Meyer, B. S. 2001, ApJ, 562, 887

Thompson, T. A., Chang, P., Quataert, E. 2004, ApJ, 611, 380

Usov, V. V. 1992, Nature, 357, 472

Uzdensky, D. A., & MacFadyen, A. I. 2006, ApJ, 647, 1192

Uzdensky, D. A. & MacFadyen, A. I., 2007, ApJ submitted (astro-ph/0609047)

van der Swaluw, E. 2003, A&A, 404, 939

Wang, L., Howell, D. A., Höflich, P., & Wheeler, J. C. 2001, ApJ, 550, 1030

Weisskopf, M. C., et al. 2000, ApJ, 536, L81

Wheeler, J. C., Yi, I., Höflich, P., & Wang, L. 2000, ApJ, 537, 810

Wilkin, F. P. 2000, ApJ, 532, 400

Woods, P. M., & Thompson, C. 2004, ArXiv Astrophysics e-prints, arXiv:astro-ph/0406133

Woosley, S. E., & Bloom, J. S. 2006, ARA&A, 44, 507

Woosley, S. E., Heger, A., & Weaver, T. A. 2002, Reviews of Modern Physics, 74, 1015